


## Deciding what to do about carpal tunnel syndrome

This short decision aid is to help you decide what to do about your carpal tunnel syndrome. You can use it on your own, or with your doctor, to help you make a decision about what's right for you at this time.

There are five main options for treating carpal tunnel syndrome. The choices are:

- **Changing lifestyle** and working habits to put less pressure on hands, and self-managing symptoms
- **Pain medications**, for quick relief from pain
- **Physical management of symptoms**, including hand exercises and wrist splints
- **Corticosteroid injections** into the carpal tunnel, to reduce pain, swelling, and pressure
- **Surgery** to reduce pressure on the carpal ligament nerve.



## What are my options?

	Lifestyle changes	Pain relief medication	Physical management	Steroid injection	Surgery
<b>What is the treatment?</b>	<p>People with carpal tunnel syndrome can choose to self-manage their condition. That just means they find out about the condition themselves, and do things to try to improve it.</p> <p>People with carpal tunnel syndrome can try making changes to their workplace or their daily lifestyle, activities, and hobbies that may help improve their symptoms.</p>	<p>Pain medications are drug treatments designed to reduce pain.</p> <p>Some of these tablets can be bought from a pharmacy without a prescription. These include paracetamol, paracetamol combined with low-strength codeine, and some non-steroidal anti-inflammatory drugs (NSAIDs) such as ibuprofen and aspirin. Other drug treatments need to be prescribed by a doctor.</p>	<p>Physical management of carpal tunnel syndrome usually means hand exercises or splinting.</p> <p>Hand exercises are exercises that involve stretching the fingers and wrist into different positions to see if this can release the pressure on the nerve. Wrist splints keep the hand and wrist straight and reduce the pressure on the trapped nerve.</p>	<p>Having a steroid injection (the full name of this type of medication is corticosteroid) means having an injection into the carpal tunnel. This is done by a doctor.</p> <p>Corticosteroids are sometimes called steroids for short, but they are not the same as the anabolic steroids used by some bodybuilders and athletes. Corticosteroids are similar to the natural chemicals the body makes to reduce swelling.</p>	<p>Surgery involves an operation on the wrist. The surgeon cuts free a band of tissue called the carpal ligament from the median nerve. The operation is usually done under local anaesthetic. This operation can be done by either open or keyhole (also called endoscopic) surgery.</p> <p>Surgery is usually only recommended for people who have severe symptoms and have tried other treatments first.</p>

	Lifestyle changes	Pain relief medication	Physical management	Steroid injection	Surgery
<b>What is the effect on pain?</b>	<p>We don't know whether lifestyle changes can reduce pain. There has not been much research in this area.</p> <p>We don't know for sure whether using specially designed keyboards or equipment can reduce pain for people with carpal tunnel syndrome. [1]</p>	<p>We don't know whether or not non-steroidal anti-inflammatory drugs (NSAIDs) or paracetamol help pain. There hasn't been much research.[2]</p>	<p>Nerve and tendon exercises may help to reduce pain. We don't know for sure because there has not been much research into this. [3]</p> <p>Wearing a wrist splint at night can help to reduce pain. About 50 in 100 people with carpal tunnel syndrome say their symptoms are improved three months after starting using a splint.[4]</p>	<p>A corticosteroid injection into the carpal tunnel can make the hand, arm, and wrist feel less painful and swollen.[5] People may feel an improvement for up to one month after the injection. [6]</p>	<p>Six months after surgery, most people say their pain has improved. [7]</p> <p>The two types of surgery (open and keyhole) seem to work as well as each other in treating the pain of carpal tunnel syndrome.[8]</p>

	Lifestyle changes	Pain relief medication	Physical management	Steroid injection	Surgery
<b>What is the effect on movement?</b>	<p>We don't know whether lifestyle changes can improve how easily people can move their hands or wrists. There has not been much research in this area.</p> <p>We don't know for sure whether using special keyboards or equipment can make it easier for people with carpal tunnel syndrome to move their hands or wrists. There has not been enough research. [9]</p>	<p>We don't know whether NSAIDs improve hand and wrist movement in carpal tunnel syndrome. There hasn't been much research.[10]</p> <p>Someone with carpal tunnel syndrome may find it easier to move their hand and wrist if they wear a splint as well as taking NSAIDs.[11]</p> <p>After six weeks of treatment, people with carpal tunnel syndrome who take NSAIDs and wear a splint say they find it easier to write and to pick up small objects. [12]</p>	<p>We don't know whether nerve and tendon exercises improve how well people can move their hands or wrists.[13]</p> <p>Wearing a wrist splint at night can help to improve how well someone with carpal tunnel syndrome can move their hand or wrist.[14]</p> <p>People who have a corticosteroid injection as well as wearing a splint find it easier to do things such as write, button clothes, grip a telephone receiver, open jars, do housework, carry grocery bags, and bathe themselves. [15]</p>	<p>Corticosteroid injections into the carpal tunnel can improve how well people can move their hands and wrists. People may feel an improvement up to one month after the injection.[16]</p> <p>Someone with carpal tunnel syndrome may find it easier to move their hand and wrist if they wear a splint as well as having a corticosteroid injection.[17]</p>	<p>Carpal tunnel surgery usually makes it easier to move the hand and wrist.[18][19][20]</p> <p>Six months after surgery, about 56 in 100 people say they can move their wrists better than before the operation.[21]</p> <p>Some people find that surgery does not improve movement or strength.[22]</p>

	Lifestyle changes	Pain relief medication	Physical management	Steroid injection	Surgery
<b>How much time is spent in hospital or having treatment?</b>	Some lifestyle changes require daily commitment, or take some getting used to, such as using new equipment. A session with an occupational therapist at work may be needed.	Some pain relief medication can be bought over the counter, without a prescription. A doctor may also prescribe pain medication. People with carpal tunnel syndrome might need to take medicines regularly to manage their pain.	<p>People may need to go to hospital to have a number of physiotherapy sessions. They won't need to stay in hospital.</p> <p>Wrist splints are usually available from larger pharmacies, or a GP may be able to recommend a suitable supplier. The splint can be used for several weeks or months, depending on how severe the symptoms are.</p>	Corticosteroid injections are usually given at a GP surgery or at a special clinic. To begin with, doctors usually recommend one injection. They may suggest more injections if necessary.[23]	<p>Surgery for carpal tunnel syndrome is done in hospital. People should be able to go home the same day.</p> <p>People who have surgery will need to wear a bandage for about two days after the operation. Their doctor may recommend gentle hand, wrist, and elbow exercises to prevent stiffness.</p> <p>It's possible that surgery will not work and a second operation will be needed.[24]</p>

	Lifestyle changes	Pain relief medication	Physical management	Steroid injection	Surgery
<b>How quickly does the treatment work?</b>	We don't know how quickly lifestyle changes or using special equipment can improve the symptoms. There has not been much research.[25]	If pain medications help, they should start working quickly, within minutes rather than hours.	<p>People who find that nerve and tendon exercises help their symptoms say that the improvement happens within three months.[26]</p> <p>People who find that wearing a wrist splint at night helps their symptoms say they start to notice an improvement after about four weeks. [27]</p>	Corticosteroid injections may start working one week after treatment.[28] People might feel an improvement for up to one month after the injection.[29]	About 80 in 100 people who have surgery say their symptoms have improved three months after the operation.[30]

	Lifestyle changes	Pain relief medication	Physical management	Steroid injection	Surgery
<b>How long do the effects of treatment last?</b>	For people who find that lifestyle changes or special equipment help, we don't know how long the benefit will last. There hasn't been much research.[31]	People need to keep taking pain medications for them to keep working. If pain medications stop helping, doctors may advise people to stop taking them.	<p>We don't know how long the benefits of nerve and tendon exercises may last, as most of the research has only looked at the first three months after treatment starts. [32]</p> <p>About 50 in 100 people say their symptoms are better three months after starting using a wrist splint.[33]</p> <p>People may continue to feel better even after they stop wearing the splint, but we don't know for certain. There hasn't been much research.[34]</p>	About 94 in 100 people say their symptoms improve soon after a corticosteroid injection. Two years after the injection this number is 60 in 100.[35]	<p>For people who benefit from surgery, the benefits last for at least six months, and can last a lifetime.[36][37]</p> <p>About 75 in 100 people who have surgery say that their symptoms improve. After two years, the number is 69 in 100.[38]</p>

	Lifestyle changes	Pain relief medication	Physical management	Steroid injection	Surgery
<b>What side effects or complications does the treatment have?</b>	<p>Making lifestyle changes, or using special keyboards or equipment, are unlikely to have any side effects or complications.[39]</p>	<p>NSAIDs can cause bleeding in the stomach when taken over a long period of time. They can also damage the kidneys. NSAIDs may increase the chance of heart attacks and strokes. NSAID tablets may not be suitable for some people with certain conditions, such as asthma, a peptic ulcer, or angina, or in people who have had a heart attack or stroke.</p> <p>Paracetamol is unlikely to cause side effects in people who take the correct dose. Taking too much can damage the liver.</p>	<p>We don't know for sure whether nerve and tendon exercises can cause unwanted effects, because the research doesn't tell us. It's possible that some exercises could make symptoms worse.[40]</p> <p>Wearing a wrist splint is generally safe. Some people may experience discomfort and swelling of the hands and wrist, difficulty falling asleep when wearing a splint at night, or pins and needles when taking it off in the morning.[41]</p>	<p>There is a chance that the nerve could be damaged when corticosteroids are injected into the carpal tunnel. The injection could also lead to an infection in the wrist.[42]</p>	<p>Carpal tunnel surgery can cause unwanted effects. The most common are skin irritation, scarring, infection, and blood collecting under the skin at the site of the wound (haematoma). [43] The scar from surgery can be irritated and painful for two to three months after the operation.</p> <p>It's possible that surgery will not work and a second operation may be needed.[44]</p>

## What are the pros and cons of each option?

People with carpal tunnel syndrome have different experiences about the health problem and views on treatment. Choosing the treatment option that is best for the patient means considering how the consequences of each treatment option will affect their life.

Here are some questions people may want to consider about treatment for carpal tunnel syndrome:

- Are they willing to wait and see what happens to their symptoms?
- Do they want to be able to do more everyday things than they can manage at present?
- Are they willing to spend time having treatment?
- Are they willing to spend time recovering from treatment?
- Are they willing to try a treatment that may take a long time to have an effect?

## How do I get support to help me make a decision that is right for me?

Go to <http://sdm.rightcare.nhs.uk/pda/carpal-tunnel-syndrome/> for more detailed information about treatments for **Carpal Tunnel Syndrome**. People using this type of information say they understand the health problem and treatment choices more clearly, and why one treatment is better for them than another. They also say they can talk more confidently about their reasons for liking or not liking an option with health professionals, friends and family.

## References

References can be viewed online at <http://sdm.rightcare.nhs.uk/pda/carpal-tunnel-syndrome/references/>